

Lepton flavor violation decays with the fourth generation neutrino ν'

Wu-Jun Huo and Tai-Fu Feng

CCAST (World Lab.), P.O. Box 8730, Beijing 100080

and

Institute of High Energy Physics, Academia Sinica, P.O. Box 918(4),

Beijing 100039, P.R. China

today

Abstract

We investigate the lepton flavor violation decays, $\tau \rightarrow \mu\gamma$, $\tau \rightarrow e\gamma$ and $\mu \rightarrow e\gamma$, in the framework of a sequential fourth generation model with a heavy fourth neutrino, ν' . Using the recent experimental bounds, we can obtain the constraints of the 4×4 leptonic mixing matrix element factors, $V_{1\nu'}$, $V_{2\nu'}$ and $V_{3\nu'}$. We find that LFV decays can give stringent constraints on the parameter space of the 4th neutrino mass $m_{\nu'}$.

At present, Standard model (SM) has to face the experimental difficulties [1, 2] which are all related to leptons. It seems to indicate the presence of new physics just around the corner will be in the leptonic part. The leptonic flavor violation decay $\tau \rightarrow \mu\gamma$ is very useful to search new physics. In this note, we consider the sequential fourth generation standard model (SM4) to investigate its contributions to some leptonic flavor violation decays, $\tau \rightarrow \mu\gamma$, $\tau \rightarrow e\gamma$ and $\mu \rightarrow e\gamma$.

From the point of phenomenology, there is a realistic question what are numbers of the fermions generation or whether there are other additional quarks or leptons. The present experiments tell us there are only three generation fermions with *light* neutrinos which mass are less smaller than $M_Z/2$ [3]. But the experiments don't exclude the existence of other additional generation, such as the fourth generation, with a *heavy* neutrino, i.e. $m_{\nu_4} \geq M_Z/2$ [4]. Many refs. have studied the SM4 [5], which is added an up-like quark t' , a down-like quark b' , a lepton τ' , and a heavy neutrino ν' in the SM. The properties of these new fermions are all the same as their corresponding counterparts of other three generations except their masses and CKM mixing, see Tab.1,

	up-like quark	down-like quark	charged lepton	neutral lepton
SM fermions	u	d	e	ν_e
	c	s	μ	ν_μ
	t	b	τ	ν_τ
new fermions	t'	b'	τ'	ν'

Table 1: The elementary particle spectrum of SM4

If there exists a very heavy fourth neutrino ν' , it can contribute to a_μ through diagram of Fig. 1. This is the electroweak interaction. Similar to that of quarks, the corresponding Lagrangian is

$$\mathcal{L} = -\frac{g}{\sqrt{2}}(\bar{\nu}'\gamma_\mu a_L V_{l\nu'}l)W^\mu + h.c. \quad (1)$$

where $a_L = (1 - \gamma_5)/2$, $V_{l\nu'}$ ($l = e, \mu, \tau$) is the (1,4), (2,4) and (3,4) elements of the

four-generation CKM matrix (4×4),

$$V_{\text{CKM}}^{\text{SM4}} = \begin{pmatrix} V_{1\nu_e} & V_{1\nu_\mu} & V_{1\nu_\tau} & V_{1\nu'} \\ V_{2\nu_e} & V_{2\nu_\mu} & V_{2\nu_\tau} & V_{2\nu'} \\ V_{3\nu_e} & V_{3\nu_\mu} & V_{3\nu_\tau} & V_{3\nu'} \\ V_{4\nu_e} & V_{4\nu_\mu} & V_{4\nu_\tau} & V_{4\nu'} \end{pmatrix} \quad (2)$$

Reverting back to the diagrams of Fig. 1, we see that the fourth neutrino contribution to decays $\tau \rightarrow e\gamma, \mu\gamma$ and $\mu \rightarrow e\gamma$ (see Fig. 1). We don't consider the usual three neutrinos' contribution because their masses are too small. For the heavy neutrino ν' , In SM4, it can induce these decays. Calculating Fig. 1, one obtains

$$\Gamma(\mu \rightarrow e\gamma) = \frac{\alpha G_F^2 m_\tau^5}{96} |V_{1\nu'} V_{2\nu'}|^2 f^2(x), \quad (3)$$

where $x \equiv m_{\nu'}^2/m_W^2$, α is the fine structure constant and

$$f(x) = \frac{-5x^3 - 5x^2 + 4x}{12(x-1)^3} + \frac{(2x^3 - x^2) \log x}{2(x-1)^4}. \quad (4)$$

Similarly, we obtain

$$\Gamma(\tau \rightarrow e\gamma) = \frac{\alpha G_F^2 m_\tau^5}{96} |V_{1\nu'} V_{3\nu'}|^2 f^2(x), \quad (5)$$

$$\Gamma(\tau \rightarrow \mu\gamma) = \frac{\alpha G_F^2 m_\mu^5}{96} |V_{2\nu'} V_{3\nu'}|^2 f^2(x). \quad (6)$$

Using the current experimental bounds of these three LFV processes [6],

$$\begin{aligned} \text{Br}(\mu \rightarrow e\gamma) &\leq 4.9 \times 10^{-11}, \\ \text{Br}(\tau \rightarrow e\gamma) &\leq 2.7 \times 10^{-6}, \\ \text{Br}(\mu \rightarrow \mu\gamma) &\leq 3.0 \times 10^{-6}. \end{aligned} \quad (7)$$

we can obtain the parameter space of other 4×4 matrix elements factors of leptonic mixing, $|V_{l\nu'} V_{l'\nu'}|^2$ (see Figs. 2, 3, 4).

From the Figs, considering the unitarity of the matrix $V_{\text{CKM}}^{\text{SM4}}$, we find the reasonable range of $m_{\nu'}^2$ is under the curve. But in our previous paper [7], the upper bounds of $m_{\nu'}$ don't exceed 80GeV from the revised $g_\nu - 2$ results. Although the low bound of $m_{\nu'}$ we get from $l \rightarrow l'\gamma$ is consistent with the present experiments [3], the upper bound of $m_{\nu'}$ seems

to conflict with the current experiments statue which there is no any new physics signals upper to hadrds GeVs. The fourth generation particles seems not to be so light. They should be several hadrad GeVs weight.

In summary, we calculate the contribution of the fourth generation to LFV decays and get an interesting result which we can exclude most values of $m_{\nu'}$. Considering the current experimental bounds, we give the parameter space of $m_{\mu'}$ and lepton mixing matrix element $V_{\mu'}$. We find that LFV processes can constrain on the neutrino mass of the fourth generation: i.e. its mass should be lighter than 80 GeV. It seems that from the lepton part, the current experiments can impose a stringent constraint on the existence of the fourth generation.

Acknowledgments


This research is supported by the the Chinese Postdoctoral Science Foundation and CAS K.C. Wong Postdoctoral Research Award Fund. We are grateful to Prof. X.M Zhang for useful discussions.

References

- [1] Y. Fukuda *et al.*, Phys. Lett. **B436** 33 (1998); Phys. Rev. Lett. **81**, 1562 (1998).
- [2] Muon g-2 Collaboration, H.N. Brown *et al.*, Phys. Rev. Lett. **86**, 2227 (2001).
- [3] G.S. Abrams et al., Mark II Collab., Phys. Rev. Lett. **63** (1989) 2173; B. Advera et al., L3 Collab., Phys. Lett. B **231** (1989) 509; I. Decamp et al., OPAL Collab., *ibid.*, **231** (1989) 519; M.Z. Akrawy et al., DELPHI Collab., *ibid.*, **231** (1989) 539; C.Caso et al., (Particle Data Group), Eur. Phys. J.C **3** (1998) 1.
- [4] Z. Berezhiani and E. Nardi, Phys. Rev. D **52** (1995) 3087; C.T. Hill, E.A. Paschos, Phys. Lett. B **241** (1990) 96.

- [5] J.F. Gunion, Douglas W. McKay, H. Pois, Phys. Lett. B **334** (1994) 339; Phys. Rev. D **51** (1995) 201; H.-J. He, N. Polonsky and S. Su, Phys.Rev. **D64** (2001) 053004.
- [6] Particle Data Group, C. Caso *et al.*, Eur. Phys. J. **C3**, 1 (1998).
- [7] W.J. Huo and F.T. Fu, hep-ph/0203173.

Figure 1: Feynmann diagram for LFV decay $\mu \rightarrow e\gamma$ induced by ν' .






Figure 2: Diagram of $|V_{1\nu'}V_{2\nu'}|^2$ to $m_{\nu'}$




Figure 3: Diagram of $|V_{1\nu'} V_{3\nu'}|^2$ to $m_{\nu'}$.




Figure 4: Diagram of $|V_{3\nu'} V_{2\nu'}|^2$ to $m_{\nu'}$.